

The Impact of Hybrid Capture-Based Comprehensive Genomic Profiling on Treatment Strategies in Patients with Solid Tumors

Umut DİŞEL^a, Fatih KÖSE^b, Ahmet BİLİCİ^c, Mustafa ÖZGÜROĞLU^d, Sezer SAĞLAM^e, Mesut ŞEKER^f, Sercan AKSOY^g, İbrahim TEK^h, Nil Molinas MANDELⁱ, Gökhan DEMİR^j, Çağatay ARSLAN^k, Mutlu DEMİRAY^l, Mehmet Akif ÖZTÜRK^m, Taflan SALEPÇİⁿ, Yeşim ERALP^o, Fatih SELÇUKBİRİCİK^p, Gökçe TEMİZAŞ^r, Ebru Gül FİDAN^r

^aClinic of Medical Oncology, Acıbadem Adana Hospital, Adana, Türkiye

^bDivision of Medical Oncology, Başkent University Adana Dr. Turgut Noyan Application and Research Center, Adana, Türkiye

^cDivision of Medical Oncology, İstanbul Medipol University Faculty of Medicine, İstanbul, Türkiye

^dDivision of Medical Oncology, İstanbul University-Cerrahpaşa, Cerrahpaşa Faculty of Medicine, İstanbul, Türkiye

^eDivision of Medical Oncology, Demiroğlu Bilim University İstanbul Florence Nightingale Hospital, İstanbul, Türkiye

^fDivision of Medical Oncology, Bezmîâlem Vakıf University Faculty of Medicine, İstanbul, Türkiye

^gDivision of Medical Oncology, Hacettepe University Institute of Cancer, Ankara, Türkiye

^hClinic of Medical Oncology, Medicana International Ankara Hospital, Ankara, Türkiye

ⁱClinic of Medical Oncology, American Hospital, İstanbul, Türkiye

^jClinic of Oncology, Acıbadem Maslak Hospital, İstanbul, Türkiye

^kClinic of Medical Oncology, Medical Park İzmir Hospital, İzmir, Türkiye

^lClinic of Medical Oncology, Medicana International İstanbul Hospital, İstanbul, Türkiye

^mClinic of Oncology, VM Medical Park Pendik Hospital, İstanbul, Türkiye

ⁿClinic of Medical Oncology, İstanbul Oncology Hospital, İstanbul, Türkiye

^oClinic of Medical Oncology, Acıbadem Maslak Hospital, İstanbul, Türkiye

^pDivision of Medical Oncology, Koç University Faculty of Medicine, İstanbul, Türkiye

^rRoche Pharma, İstanbul, Türkiye

ABSTRACT Objective: The development of bioinformatics and comprehensive genomic profiling (CGP) has provided insights into the applicability and functionality of the genomic alterations (GA). In this study, we evaluated the impact of CGP on the treatment plan and outcomes in a significant number of patients. **Material and Methods:** We carried out a retrospective case-control study on 164 adult patients with advanced solid tumors from 15 oncology centers in Türkiye. **Results:** In all cases, CGP was performed within 23.8 [standard deviation (SD)±32.1] months of initial diagnosis. Non-small cell lung carcinoma, breast cancer, unknown primary carcinoma, colorectal carcinoma, and sarcoma were among the most common tumor types, accounting for 61.5% of all cases. CGP was performed immediately after the diagnosis of advanced cancer in 13 patients (7.9%). In 158 patients (96.4%), at least one GA was found as per the CGP report. Also, in the reports, the average tumor mutational burden (TMB) and GAs were 7.3 (SD±8.7) mut/Mb and 3.5 (SD±2.0), respectively. According to CGP reports, 58 patients had 79 evidence-based drug suggestions for their particular tumor type, whereas 97 patients had 153 evidence-based drug suggestions for another tumor type. After the primary oncologist interpreted the CGP reports, significant changes were made to the treatment of 35 (21.3%) patients. **Conclusion:** We strongly believe that in the future, high-TMB or other tumor-agnostic biomarkers will become much more affordable, and CGP will serve as one of the major decision-making tools for the treatment of patients along with pathological, radiological or laboratory tests.

Keywords: Comprehensive genomic hybridization; genomics; neoplasms; retrospective studies

In recent decades, considerable progress has been made in understanding the molecular pathways of tumor progression and development.¹ Treatment strategies for major cancer types have been trans-

formed by the identification of a key mutation that is known as “driver mutation” present in a molecular pathway.² For instance, anaplastic lymphoma kinase (ALK) and epidermal growth factor receptor (EGFR)

Correspondence: Umut DİŞEL

Clinic of Medical Oncology, Acıbadem Adana Hospital, Adana, Türkiye

E-mail: doctorhope@gmail.com

Peer review under responsibility of Journal of Oncological Sciences.

Received: 01 Mar 2022 **Accepted:** 13 Jun 2022 **Available online:** 01 Jul 2022

2452-3364 / Copyright © 2022 by Turkish Society of Medical Oncology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



in lung cancer or B-RAF inhibition in melanoma have significantly changed our way to the treatment of these tumors.³ This has led to a rising interest in precision medicine and has made it increasingly evident to most physicians that interventions in oncology have shifted from histology-based treatment protocols to histology-agnostic treatment protocols based on the driver mutations.⁴ However, so far, the US Food and Drug Administration (FDA) has approved tumor agnostic treatments only for patients with tumor mutational burden-high (TMB-H), microsatellite instability-high, and neurotrophic tyrosine receptor kinase fusion-positive tumours.⁵

Recently, Comprehensive Genomic Profiling (CGP) has become more popular among oncology practitioners. CGP is a next-generation sequencing (NGS) approach that detects novel and known variants for the four main classes of genomic alterations (GA) and signatures. It provides prognostic, diagnostic, and predictive insights for all cancer types, which in turn aids in personalized research or treatment decisions for patients.⁵ Moreover, in oncology, master protocols (basket, umbrella trials with or without adaptive clinical trials) aiming to compare the effectiveness of matched treatment with standard treatment protocols are encouraged by the FDA, National Cancer Institute, and European Society for Medical Oncology (ESMO).⁶ In 2019, there were 89 basket trials on a wide range of tumor types to define the role of tumor-agnostic molecular-based treatment approaches.⁶ However, the integration of CGP with molecularly guided treatment into the routine oncology practice still poses a challenge due to several practical and financial limitations. In literature, there is scarce data available on how CGP results impact and change the treatment approach in cancer patients.^{2,7} Therefore, we aimed to evaluate the non-CGP/hotspot testing and to identify the GAs and matched therapies or suitable clinical trials based on the hybrid capture-based CGP report. We also investigated the impact of CGP on the treatment plan in a significant number of patients with advanced-stage solid tumors.

MATERIAL AND METHODS

The research was designed as a multicenter, descriptive (observational), and retrospective cohort study.

All patients with advanced solid tumors whose CGP reports were reviewed by a physician prior to enrollment and who were followed up and treated between January 2015 and December 2019 at 15 participating centers were included. Patients, who did not provide informed consent for the use of their medical data, were excluded (except for the patients who were deceased). Statistical analysis included 164 patients who met all inclusion criteria. The study was approved by the Ethical Review Committee of the Cerrahpaşa Faculty of Medicine at İstanbul University (date: June 18, 2019, no: C-03) and was carried out in accordance with the Declaration of Helsinki and Good Clinical Practice guidelines.

Medical records of eligible patients were reviewed on admission or first presentation for their type of cancer, lines of previous chemotherapy, number of prior treatments, the result of hybrid capture-based CGP, the number of actionable mutations defined by hybrid capture-based CGP, list of approved drugs for diseases, and potential eligibility for any clinical trials worldwide. In addition, the treatments used after genomic profiling were also listed.

For statistical analysis, descriptive methods were used, with numerical variables being reported as mean with standard deviation (SD) or median with interquartile range (IQR) and categorical variables as the frequency with percentage.

RESULTS

In total, 166 patients with a histologically confirmed diagnosis of malignant tumor were found eligible for the study. Statistical analysis included 164 patients who met all the inclusion criteria. 114 (69.5%) patients died before the time of data analysis. Among all patients, 82 (50%) were male, and the mean age was 55.3 (SD±14.4) years. The overall clinical and demographic characteristics of the patients are presented in [Table 1](#). The patients with a diagnosis of non-small-cell lung carcinoma (NSCLC), breast cancer, carcinoma of unknown primary (CUP), colorectal carcinoma and sarcoma represented 61.5% (n=101) of all cases, including 33 (20.1%), 25 (15.2%), 17 (10.4%), 14 (8.5%), and 12 (7.3%) patients, respectively ([Table 2](#)).

TABLE 1: Demographics and baseline characteristics.

		Total n (%)	NSCLC	Breast	Unknown		Colorectal carcinoma	Sarcoma	Pancreatic	Stomach	Other
					primary carcinoma						
Gender	Male	82 (50)	23 (69.7%)	0 (0.0%)	9 (52.9%)	9 (64.3%)	7 (58.3%)	7 (77.8%)	6 (66.7%)	21 (46.7%)	
	Female	82 (50)	10 (30.3%)	25 (100.0%)	8 (47.1%)	5 (35.7%)	5 (41.7%)	2 (22.2%)	3 (33.3%)	24 (53.3%)	
Age	Median (IQR)	55.0 (45-66)	61.0 (52-65)	51.0 (42.5-58)	56.0 (46.5-63.5)	53.0 (47-65)	43.0 (31-66.8)	70.0 (53-75.5)	59.0 (44-66)	54.0 (40-6)	
	Mean (SD)	55.3±14.4	59.5±11.1	51.0±10.6	56.5±13.4	54.6±11.9	46.8±18.2	65.8±13.6	55.1±14.7	54.4±16.8	
Patient status	Alive	50 (30.5)	8 (24.2%)	6 (24.0%)	6 (35.3%)	3 (21.4%)	5 (41.7%)	1 (11.1%)	2 (22.2%)	19 (42.2%)	
	Dead	114 (69.5)	25 (75.8%)	19 (76.0%)	11 (64.7%)	11 (78.6%)	7 (58.3%)	8 (88.9%)	7 (77.8%)	26 (57.8%)	

NSCLC: Non-small-cell lung carcinoma; IQR: Interquartile range; SD: Standard deviation.

TABLE 2: Histology of tumors (n=164).

Diagnosis	Frequency	Percent
NSCLC	33	20.1
Breast	25	15.2
Unknown primary carcinoma	17	10.4
Colorectal	14	8.5
Sarcoma	12	7.3
Pancreatic	9	5.5
Stomach	9	5.5
Others	45	27.4
Total	164	100.0

Others: Ovary, hepatobiliary carcinoma, head and neck, melanoma, uterus, brain, neuroendocrine, prostate, SCLC, bladder carcinoma, cervix squamous cell carcinoma, epidermoid carcinoma, kidney, larynx, nasopharynx, and paranasal, small intestine, testis; NSCLC: Non-small-cell lung carcinoma.

In total, 44 positive GAs were detected by local genomic testing [polymerase chain reaction, fluorescence in situ hybridization (FISH)-chromogenic in situ hybridization, and hotspot gene paneling] in the study group (Table 3).

TABLE 3: Positive genomic markers detected by classical diagnosis methods.

	n	Percent (in 164 patients)
ALK	3	1.8
BRAF	7	4.3
EGFR	8	4.9
PDL1	4	2.4
ROS	2	1.2
HER2 FISH	5	3.0
KRAS	9	5.5
MSI	2	1.2
Other	4	2.4
Total	44	26.8

Some patients had more than one record; ALK: Anaplastic lymphoma kinase; EGFR: Epidermal growth factor receptor; FISH: Fluorescence in situ; MSI: Microsatellite instability-high.

CGP test was performed with a median of 13.7 (IQR, 4.4-30.4) months following the diagnosis of advanced cancer. It was performed at baseline in 13 patients (7.9%) and after the first-, second-and third-line of systemic therapy in 72 (43.9%), 24 (14.6%), and 55 (33.5%) patients, respectively.

Figure 1 shows the frequencies of GAs detected by CGP according to tumor types. The most common GAs were TP53, KRAS, CDKN2A/B, PIK3CA, and MYC. According to the CGP report, at least one GA was found in 158 (96.4%) patients. In total, 633 gene alterations were detected among 164 patients. The mean number of alterations per patient was 3.5 (SD±2.0) [median 3.5 (IQR 2.0-5.0)].

Among all patients, the mean TMB was 7.3 (SD±8.7) mut/Mb. However, it was higher than 10 mut/MB in 32 (22.4%) patients, which was also clinically relevant. In NSCLC patients, TMB≥10 10 mut/MB was 53.8%, which was substantially higher compared to other tumor types, although the mean TMB in NSCLC was 10.4 (SD±7.1) mut/Mb (Table 4). However, all the patients were microsatellite stable.

CGP reports of the patients analyzed showed that 58 patients had 79 evidence-based drug suggestions for their tumor type, whereas 97 patients had 153 evidence-based drug suggestions for another tumor type. Moreover, at the time of the study, 126 (76.8%) patients were potentially eligible for 267 actively recruiting clinical trials.

The treatment strategies used after the CGP test can be classified as chemotherapy, targeted therapy, and immunotherapy with a rate of 79.1%, 15.2%, and 15.9%, respectively. After interpretation of the CGP




FIGURE 1: The distribution of genomic alterations according to tumor types (genes altered in a single sample are not shown). NSCLC: Non-small-cell lung carcinoma.

Diagnosis	TMB \geq 10	
	n	%
NSCL (n=26)	14	53.8
Breast (n=24)	2	8.3
Unknown primary (n=15)	2	13.3
Colorectal (n=13)	2	15.4
Sarcoma (n=12)	1	8.3
Pancreatic (n=5)	1	20.0
Stomach (n=8)	3	37.5
Other (n=40)	7	17.5
Total (n=143)	32	22.4

TMB: Tumor mutational burden; NSCLC: Non-small-cell lung carcinoma.

by the primary oncologist, substantial changes were made in the treatment decisions of 35 (21.3%) patients. We found that the chemotherapy had a proportion of 88.8% in the treatment lines prior to CGP, and it was reduced to 79.1% following CGP, while the proportion of targeted therapy increased from 11.3% to 15.2% and immunotherapy from 4.2% to 15.9%. A Sankey diagram in Figure 2 shows the change in treatment regimens after the CGP test.

DISCUSSION

A study by De Falco et al. reported that CGP was performed at baseline and after the first-line treatment in 20% and 50% of the patients, respectively.⁴ In our study, CGP was performed in 7.9% and 43.9% of the patients at baseline and during second-line treatment, respectively. It appears that physicians in Türkiye perform CGP less frequently at the baseline com-

pared to their Italian colleagues, which can be attributed to several factors. First, the samples were sent at the discretion of the patient’s primary physician, and thus the results could not be representative of the general population. Second, in Türkiye, the hotspot gene NGS is more accessible and affordable than CGP. Therefore, we believe that most of the clinicians in Türkiye first try to perform local NGS with hotspot gene panel and then perform the CGP in patients with negative or inconclusive outcomes. On the contrary, these results also showed high motivation of the medical oncologist in Türkiye to find an effective druggable target prior to the second-line cytotoxic chemotherapy.

In our study, at least one GA was found in 158 (96.4%) patients with a median of 3.5 (IQR 2.0-5.0) GA per patient. The studies by Wheler et al. and Frampton et al. reported that it was 5 and 3.06 (0-23), respectively.^{8,9} In addition, multiple studies with patients with different tumors have also reported detection of at least one GA in 82.1-93.5% of the cases by CGP test.^{4,8,10} The most common GAs detected by CGP were TP53 (46.3%), KRAS (18.3%), CDKN2A/B (15.9%), PIK3CA (14.6%), and MYC (13.4%). Unfortunately, most of these mutations are not druggable targets compared to EGFR, ALK, or BRAF mutations. Additionally, Fumagalli et al. reported that when they performed CGP in patients with NSCLC, they found at least one GA in 87% of their 76 patients.¹¹ In our study, although targeted mutations were detected in 31/164 (18.9%) patients [EGFR (4.9%), BRAF (4.3%), KRAS (3.0%), PDL1




FIGURE 2: Treatment regimens before and after the CGP testing.
 CT: Chemotherapy; CGP: Comprehensive genomic profiling.

(2.4%), HER2 FISH (2.4%), and ALK (1.8%)] using classical methods, a further genomic examination was required in these patients. For this reason, it would be more reasonable to conduct CGP for the patients at the baseline.¹² In a report of ESMO Precision Medicine Working Group, which presents its recommendations on the use of NGS for patients with metastatic cancer, it is recommended to use tumor multigene NGS, especially in patients presenting with advanced NSCLC, prostate, ovarian cancers, and cholangiocarcinoma.¹²

Our genomic profiling results revealed 79 evidence-based drug suggestions for the patients' tumor type and 153 evidence-based drug suggestions for another tumor type in 58 (35.4%) and 97 (59.1%) patients, respectively. After interpretation of the CGP by the primary oncologist, significant changes were made in treatment strategies in 35 (21.3%) patients. Drilon et al. reported that GA leads to major treatment changes in 26% of the patients, which was compatible with the National Comprehensive Cancer Network (NCCN) guideline recommendations.¹³ In the literature, major treatment changes were reported in 21-45% of the cases after CGP.^{7,14} In our study, the

proportion of cytotoxic chemotherapy administered prior to CGP was 88.8%; this percentage decreased to 79.1% in the treatments given after CGP, whereas the proportion of targeted therapy increased from 11.3% to 15.2%. We strongly believe that this change is associated with increased detection of targetable driver mutations by CGP.

The overall percentage of patients with TMB \geq 10 muts/Mb was 22.4%, compared to 53.8% in NSCLC patients. Our results showed that TMB \geq 10 muts/Mb was detected in patients with gastric cancer, CUP, and sarcoma, with a mean value of 9.6 (SD \pm 5.9) 7.5 (SD \pm 15.3), and 7.3 (SD \pm 11.8) muts/Mb, respectively. These tumors showed higher TMB compared to other histology in the study. TMB is a new biomarker for cancer immunotherapy. Since TMB has a positive correlation with the neoantigen load, it has been reported that high TMB is associated with increased anti-tumor immune responses and greater utilization of immune checkpoint inhibitors.¹⁴ This relationship between TMB and immune-checkpoint inhibitors has led to the first tumor-agnostic drug approval by the FDA.¹⁵ NCCN strongly recommends that each cancer patient with good performance sta-

tus and a lack of standard treatment options should undergo TMB testing to seek potential benefits from checkpoint inhibitors.¹⁶

Multiple studies using various profiling approaches have shown that molecular profiling-guided therapy may have significant benefits in terms of response rates and survival rates. Some studies used a single analytical approach (i.e., NGS), while others applied multiple analytical methods for extensive molecular profiling. Multiplatform profiling analyses could find more druggable molecular targets and guide a higher proportion of patients toward matched-treatment strategies.¹⁷ In our study, due to the lack of effectiveness of data in our retrospective analysis, we could not reach a firm conclusion on the response rate and survival rate of our patients.

Our results provide conclusive evidence for the effectiveness of CGP in daily oncology practice using a relatively high number of patients in full datasets from Türkiye, which is a developing country. However, a few limitations are worth noting. First, patients were selected at the discretion of the patient's primary oncologist; thus, the results cannot be generalized. Second, the retrospective nature of the study without any comparison of CGP-based treatment with standard care in effectiveness parameters (response rate and survival rate) precludes us from drawing general conclusions. Third, there was no formal molecular tumor board to evaluate the result of the CGP and all treatment suggestions by the primary oncologist. Another limitation of the study is that there was no direct comparison with hot spot or single mutation testing, and we believe that it would be an interesting area for further research. On the other hand, our study had the strength to examine an extensive number of alterations at the same time using CGP.

CONCLUSION

In conclusion, our data showed that CGP identified GA in over 90% of cancer patients. Bioinformatics evaluation of CGP resulted in evidence-based drug options in patients' tumor type and another tumor

type for 58 (35.4%) and 97 (59.1%) patients, respectively. Moreover, our study revealed that the treatment strategies changed after CGP in 35 (21.3%) patients. However, due to reimbursement issues in Türkiye, we could not use pembrolizumab in TMB-H patients. But we strongly believe that the affordability of the high-TMB or other tumor-agnostic drugs will significantly increase in the future, and CGP testing will serve as one of the major decision-making tools for the patients along with pathological, radiological, or laboratory tests.

Source of Finance

During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject, nor from a company that provides or produces medical instruments and materials which may negatively affect the evaluation process of this study.

Conflict of Interest

No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

Authorship Contributions

Idea/Concept: Umut Dişel, Fatih Köse, Ahmet Bilici, Mustafa Özgüroğlu, Sezer Sağlam, Mesut Şeker, Sercan Aksoy, İbrahim Tek, Yeşim Eralp, Nil Molinas Mandel, Gökhan Demir, Çağatay Arslan, Mutu Demiray, Taflan Salepci, Mehmet Akif Öztürk, Fatih Selçukbiricik, Gökçe Temizaş, Ebru Gül Fidan; **Design:** Umut Dişel, Fatih Köse, Ahmet Bilici, Mustafa Özgüroğlu, Sezer Sağlam; **Control/Supervision:** Umut Dişel, Fatih Köse, Ahmet Bilici; **Data Collection and/or Processing:** Umut Dişel, Fatih Köse, Ahmet Bilici, Mustafa Özgüroğlu, Sezer Sağlam, Mesut Şeker, Sercan Aksoy, İbrahim Tek, Yeşim Eralp, Nil Molinas Mandel, Gökhan Demir, Çağatay Arslan, Mutlu Demiray, Taflan Salepci, Mehmet Akif Öztürk, Fatih Selçukbiricik; **Analysis and/or Interpretation:** Umut Dişel, Fatih Köse, Ahmet Bilici, Mustafa Özgüroğlu, Sezer Sağlam; **Literature Review:** Umut Dişel, Fatih Köse, Ahmet Bilici, Mustafa Özgüroğlu, Sezer Sağlam; **Writing the Article:** Umut Dişel, Fatih Köse, Ahmet Bilici, Mustafa Özgüroğlu, Gökçe Temizaş, Ebru Gül Fidan; **Critical Review:** Umut Dişel, Fatih Köse, Ahmet Bilici, Mustafa Özgüroğlu, Gökçe Temizaş, Ebru Gül Fidan; **References and Fundings:** Gökçe Temizaş, Ebru Gül Fidan.

REFERENCES

- Johnson TM. Perspective on precision medicine in oncology. *Pharmacotherapy*. 2017;37(9):988-989. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Cheng DT, Mitchell TN, Zehir A, et al. Memorial sloan kettering-integrated mutation profiling of actionable cancer targets (MSK-IMPACT): a hybridization capture-based next-generation sequencing clinical assay for solid tumor molecular oncology. *J Mol Diagn*. 2015;17(3):251-264. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Schwaederle M, Zhao M, Lee JJ, et al. Impact of precision medicine in diverse cancers: a meta-analysis of phase II clinical trials. *J Clin Oncol*. 2015;33(32):3817-3825. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- De Falco V, Poliero L, Vitello PP, et al. Feasibility of next-generation sequencing in clinical practice: results of a pilot study in the Department of Precision Medicine at the University of Campania 'Luigi Vanvitelli'. *ESMO Open*. 2020;5(2):e000675. Erratum in: *ESMO Open*. 2020;5(3). [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Looney AM, Nawaz K, Webster RM. Tumour-agnostic therapies. *Nat Rev Drug Discov*. 2020;19(6):383-384. [[Crossref](#)] [[PubMed](#)]
- Singh AP, Shum E, Rajdev L, et al. Impact and diagnostic gaps of comprehensive genomic profiling in real-world clinical practice. *Cancers (Basel)*. 2020;12(5):1156. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Tsimberidou AM, Hong DS, Ye Y, et al. Initiative for molecular profiling and advanced cancer therapy (IMPACT): an MD anderson precision medicine study. *JCO Precis Oncol*. 2017;2017:PO.17.00002. [[PubMed](#)] [[PMC](#)]
- Wheler JJ, Janku F, Naing A, et al. Cancer therapy directed by comprehensive genomic profiling: a single center study. *Cancer Res*. 2016;76(13):3690-3701. [[Crossref](#)] [[PubMed](#)]
- Frampton GM, Fichtenholtz A, Otto GA, et al. Development and validation of a clinical cancer genomic profiling test based on massively parallel DNA sequencing. *Nat Biotechnol*. 2013;31(11):1023-1031. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Sailer V, Eng KW, Zhang T, et al. Integrative molecular analysis of patients with advanced and metastatic cancer. *JCO Precis Oncol*. 2019;3:PO.19.00047. [[PubMed](#)] [[PMC](#)]
- Fumagalli C, Catania C, Ranghiero A, et al. Molecular profile of advanced non-small cell lung cancers in octogenarians: the door to precision medicine in elderly patients. *J Clin Med*. 2019;8(1):112. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Mosele F, Remon J, Mateo J, et al. Recommendations for the use of next-generation sequencing (NGS) for patients with metastatic cancers: a report from the ESMO Precision Medicine Working Group. *Ann Oncol*. 2020;31(11):1491-1505. [[Crossref](#)] [[PubMed](#)]
- Drilon A, Wang L, Arcila ME, et al. Broad, hybrid capture-based next-generation sequencing identifies actionable genomic alterations in lung adenocarcinomas otherwise negative for such alterations by other genomic testing approaches. *Clin Cancer Res*. 2015;21(16):3631-3639. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Massard C, Michiels S, Fertié C, et al. High-throughput genomics and clinical outcome in hard-to-treat advanced cancers: results of the MOSCATO 01 Trial. *Cancer Discov*. 2017;7(6):586-595. [[Crossref](#)] [[PubMed](#)]
- Marabelle A, Fakih MG, Lopez J, et al. Association of tumour mutational burden with outcomes in patients with select advanced solid tumours treated with pembrolizumab in KEYNOTE-158. *Ann Oncol*. 2019;30(5):477-478. [[Crossref](#)]
- Chang H, Sasson A, Srinivasan S, et al. Bioinformatic methods and bridging of assay results for reliable tumor mutational burden assessment in non-small-cell lung cancer. *Mol Diagn Ther*. 2019;23(4):507-520. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]
- Zimmer K, Kocher F, Spizzo G, Salem M, Gastl G, Seeber A. Treatment according to molecular profiling in relapsed/refractory cancer patients: a review focusing on latest profiling studies. *Comput Struct Biotechnol J*. Mar 2019;17:447-453. [[Crossref](#)] [[PubMed](#)] [[PMC](#)]